

Inferring soil moisture memory from streamflow observations using a simple water balance model

Rene Orth¹, Randal D. Koster² and Sonia I. Seneviratne¹

DRAFT MANUSCRIPT

20 June 2013

¹Institute for Atmospheric and Climate Science, ETH Zurich, Universitätstrasse 16, CH-8092 Zurich, Switzerland (rene.orth@env.ethz.ch, sonia.seneviratne@env.ethz.ch)

²Global Modeling and Assimilation Office, NASA Goddard Space Flight Center, Greenbelt, MD 20771 (randal.d.koster@nasa.gov)

Abstract

Soil moisture is known for its integrative behavior and resulting memory characteristics. Soil moisture anomalies can persist for weeks or even months into the future, making initial soil moisture a potentially important contributor to skill in weather forecasting. A major difficulty when investigating soil moisture and its memory using observations is the sparse availability of long-term measurements and their limited spatial representativeness. In contrast, there is an abundance of long-term streamflow measurements for catchments of various sizes across the world. We investigate in this study whether such streamflow measurements can be used to infer and characterize soil moisture memory in respective catchments. Our approach uses a simple water balance model in which evapotranspiration and runoff ratios are expressed as simple functions of soil moisture; optimized functions for the model are determined using streamflow observations, and the optimized model in turn provides information on soil moisture memory on the catchment scale. The validity of the approach is demonstrated with data from three heavily monitored catchments. The approach is then applied to streamflow data in several small catchments across Switzerland to obtain a spatially distributed description of soil moisture memory and to show how memory varies, for example, with altitude and topography.

1 Introduction

Among the variables of the climate system, soil moisture has potentially important memory (persistence) characteristics. If soil moisture anomalies, as induced by precipitation anomalies, persist into subsequent weeks, and if these long-lasting anomalies are then translated to the atmosphere through their impacts on the surface energy balance, soil moisture memory may have profound implications for climate variability and prediction.

The role of soil moisture memory in climate, however, is still not completely understood. Complexity arises, for example, from the fact that while a soil moisture persistence signal can be translated to the atmosphere through evaporation anomalies (i.e., through soil moisture-evapotranspiration coupling and land-atmosphere interactions), these evaporation anomalies in turn act to reduce any original soil moisture anomaly; that is, a soil moisture anomaly, when it affects the surface fluxes, also acts to limit its own lifetime (although positive feedbacks with precipitation could also enhance it, e.g. *Koster and Suarez* 2001). In considering this balancing act, it is instructive to consider two competing and extreme scenarios. In the first scenario, evaporation processes annihilate a soil moisture anomaly within a day or two of its formation; soil moisture memory would then be small, and its effects on climate variability would necessarily be minimal. In the second scenario, the soil moisture anomaly does not affect evaporation or runoff and thereby persists indefinitely; here again, because the atmosphere or rivers cannot feel the anomaly, impacts on climate variability would necessarily be small. Evidence exists to show that neither of these extremes wholly captures the way nature works. In many regions, an important middle ground is achieved: soil moisture anomalies have been observed to persist for weeks to months (*Vinnikov and Yesserkepova* 1990, *Entin et al.* 2000, *Seneviratne et al.* 2006), and their impacts on atmospheric variability do indeed manifest themselves at those timescales, as demonstrated by various studies which quantify the impact of soil moisture initialization on the skill of sub-seasonal precipitation and/or temperature forecasts (e.g., *Viterbo and Betts* 1999, *Koster et al.*

2004, *Douville* 2010, *Koster et al.* 2010b, *Seneviratne et al.* 2010) or identified lag correlations between surface moisture deficits and temperature extremes (e.g., *Hirschi et al.* 2011, *Mueller and Seneviratne* 2012). This memory, at the same time, also allows soil moisture initialization to contribute significant skill to seasonal streamflow forecasts (e.g., *Koster et al.* 2010a, *Mahanama et al.* 2012).

The existence of this useful middle ground makes soil moisture memory worthy of careful study. A critical step in this understanding is the characterization of memory and its variations across the globe. Unfortunately, such a characterization is not straightforward. A major obstacle is the limited availability of long-term soil moisture measurements (e.g. *Robock et al.* 2000, *Seneviratne et al.* 2010, *Dorigo et al.* 2011). Ground measurements of soil moisture are only available at the point scale, which implies some limitation in their spatial representativeness. Although spatial variability should not be overstated (e.g. *Mittelbach and Seneviratne* 2012), different hydrological dynamics may be active, for example, over adjacent grassland and forest areas (*Teuling et al.* 2010b, *Orth and Seneviratne* 2012a). Also, model estimates of soil moisture cannot be used for persistence studies, given the dependence of simulated soil moisture persistence on generally unvalidated model assumptions.

In contrast, streamflow measurements are widely available, they generally cover longer periods, and they represent an integral of hydrological processes over an area. Because streamflow itself responds to soil moisture variations (see also *Kirchner* 2009 and *Mahanama et al.* 2012), it is natural to ask whether streamflow measurements contain useful information on catchment-scale soil moisture anomalies and soil moisture memory. We address this question in this paper. Using an adaptation of the simple water-balance model of *Koster and Mahanama* (2012), streamflow measurements are translated into fitted functional relationships between soil moisture and both runoff and evapotranspiration. These fitted relationships in turn provide estimates of soil moisture memory. The approach is successfully validated in three heavily monitored catchments in central

Europe and is then applied to several near-natural catchments in Switzerland, providing a spatial picture of how soil moisture memory varies across the country. The analysis shows how soil moisture memory is affected by both, geomorphological controls (e.g., altitude, topography, and catchment size) as well as meteorological controls (e.g., dryness index and the potential for externally-induced memory from the atmospheric forcing to be transmitted into the soil).

2 Methodology

2.1 Simple Water-balance Model

Koster and Mahanama (2012) (hereafter referred to as KM12) developed a simple water-balance model to study the influence of soil moisture on hydroclimatic means and variability on large spatial and temporal scales. We use a similar approach in the present study. However, because we focus here on soil moisture memory in small catchments on daily to weekly time scales, we introduce several new features to the model, as described below.

2.1.1 Water Balance Equation

As in KM12, the model used here is based on the following water balance equation:

$$w_{n+\Delta t} = w_n + (P_n - E_n - Q_n) \Delta t \quad (1)$$

where w_n denotes the model's sole prognostic variable: the total soil moisture content (in mm) at time step n . The value of w_n is altered by precipitation P_n , evapotranspiration E_n , and runoff Q_n (all in $\frac{mm}{d}$) accumulated from time step n to $n + \Delta t$ to yield the soil moisture at the next time step, $w_{n+\Delta t}$. As in KM12, we run the model here with a time step of one day ($\Delta t = 1d$).

2.1.2 Evapotranspiration

As in KM12, we assume simple dependencies of evapotranspiration (normalized by net radiation (in $\frac{W}{m^2}$)) and runoff (normalized by precipitation) on soil moisture. We use the following equation to capture the control of soil moisture on the ratio of evapotranspiration to net radiation, or ET ratio:

$$\frac{\lambda \rho_w E_n}{R_n} = \beta_0 \left(\frac{w_n}{c_s} \right)^\gamma \text{ with } \gamma > 0 \text{ and } \beta_0 \leq 1 \quad (2)$$

where λ is the latent heat of vaporization (in $\frac{J}{kg}$) and ρ_w is the density of water (in $\frac{kg}{m^3}$). Soil moisture is scaled by the soil water holding capacity c_s (in mm) so that the function operates on the degree of saturation (unitless). The unitless exponent γ ensures that the function is strictly monotonically increasing, so that the ET ratio increases with soil moisture. The factor β_0 (also unitless) reflects the residual plant and soil evaporative resistance under conditions which are not soil moisture-limited (e.g. *Seneviratne et al.* 2010). This factor therefore prevents the complete conversion of available net radiation into ET even when water is fully available (reflecting, for example, the fact that even with no water stress, transpiring water must still travel through the vegetation).

2.1.3 Runoff and streamflow

Even if runoff in nature is controlled by many variables, we assume that it depends on precipitation and soil moisture only, according to the equation:

$$\frac{Q_n}{P_n} = \left(\frac{w_n}{c_s} \right)^\alpha \text{ with } \alpha \geq 0 \quad (3)$$

As with the exponent γ in (2), the unitless exponent α ensures that the runoff ratio $\frac{Q_n}{P_n}$ increases

monotonically with soil moisture. Note that runoff as defined here (which includes, in effect, both overland flow and drainage to baseflow-producing groundwater) is distinct from streamflow, as measured at a stream gauge site; the latter quantity includes delays associated with the subsurface water transport to the streambeds and the transport of the surface water to the stream gauge site. Based on sensitivity tests, we found that accounting explicitly for this distinction between runoff and streamflow improves the model's performance in comparison to the KM12 version (not shown). We thus compute streamflow from the simulated runoff values by imposing a delay characterized by a timescale τ :

$$S_{n+t} = Q_n \frac{1}{\tau} e^{-\frac{t}{\tau}} \quad (4)$$

where the streamflow S_{n+t} corresponds to the streamflow produced at time $n+t$ associated with the surface runoff formed at time n . The integral of $\frac{1}{\tau} e^{-\frac{t}{\tau}}$ as $t \rightarrow \infty$ equals 1, ensuring that the full complement of assumed runoff water (i.e., $P_n \left(\frac{w_n}{c_s}\right)^\alpha$, from Equation (3)) does contribute to streamflow at some time. The parameter $\frac{1}{\tau}$ determines how quickly the runoff is transformed into streamflow, whereas τ corresponds to the recession time scale, expressed in days. Using Equation (4), the streamflow accumulated over the m -th time step after the precipitation event is:

$$S_{n+m\Delta t} = \left(\frac{w_n}{c_s}\right)^\alpha P_n \int_m^{m+\Delta t} \frac{1}{\tau} e^{-\frac{t}{\tau}} dt = Q_n \left(e^{-\frac{m\Delta t}{\tau}} - e^{-\frac{(m+1)\Delta t}{\tau}} \right) \quad (5)$$

With this equation we can express the streamflow at any time step as the accumulation of the effects of all runoff amounts generated during the preceding 60 time steps:

$$S_n = \sum_{i=0}^{60} Q_{n-i\Delta t} \left(e^{-\frac{i\Delta t}{\tau}} - e^{-\frac{(i+1)\Delta t}{\tau}} \right) \quad (6)$$

Note that in order to make sure that all the generated runoff is transformed into streamflow, we would in principle need to use an infinite number of time steps. Sixty time steps is an arbitrary but tractable number that allows us to account for 99% or more of the runoff water.

2.1.4 Model Integration

Assuming that values for the five parameters in Equations (2) and (6) (namely, c_s , β_0 , γ , α , and τ) can be determined, Equation (1) can be driven with daily values of precipitation and net radiation over any time period of interest to produce daily time series of total soil moisture, w_n , as well as daily time series of runoff and ET. In contrast to KM12, who used monthly precipitation observations (equally distributed across the days of a given month) and an observed seasonal climatology of net radiation to force their model, we employ daily observations of precipitation and radiation. Unlike KM12, we do not include a snow layer in the model as our study focuses on the growing season.

In fact, due to the limitation of using a daily (rather than a finer) time step, we integrate instead an implicit form of Equation (1), a form that effectively computes the evaporation and runoff for a given day based on the soil moisture content at the end of that day:

$$w_{n+\Delta t} - w_n = P_n - E_{n+\Delta t} - Q_{n+\Delta t} \approx \frac{P_n - E_n - Q_n}{1 + E'_n + Q'_n} \quad (7)$$

where the prime ($'$) indicates the derivative with respect to soil moisture, evaluated at w_n . Note that even with this correction, the time-discretized equation is still not perfectly solved because the functions $E(w)$ and $Q(w)$ are not linear but (partly strongly) curved.

Running the model requires the initialization of the soil moisture prognostic variable. We spin-up the model by integrating it over five years prior to the start of a simulation.

2.2 Optimization of streamflow, runoff and evapotranspiration parameters

We optimize the above model with daily data from 16 European catchments, three of which have been previously examined in *Orth and Seneviratne* (2012a) (hereafter referred to as OS12). We use precipitation and radiation observations in these catchments to force the model. We then identify, separately for each catchment, the optimal set of values for the 5 parameters in Equations (2) and (6), that is, the set of values that allows the modeled streamflow S_n (Equation (6)) to agree most closely with observed streamflow.

The accuracy of the modeled streamflow is measured with a time correlation against observed streamflow. The correlation period is limited to July through September to avoid any impact of snow, which is not included in the model (May-September for warmer site San Rossore). The absence of snow is supported by daily average temperatures that are always above 0 degrees Celsius during the correlation period. Note that while applied here to specific basins in Switzerland, the simple water balance model is generally applicable to any region and time period where streamflow is present.

One way to find the optimal set of values for the five parameters at each catchment would be to run the model using all possible combinations of values. Capturing the optimal values in this way with some accuracy, however, would be computationally prohibitive. To work around this problem, we developed an alternative procedure (see Appendix A) to reduce the number of model runs required to yield a reliable optimal parameter set (see Table 1).

2.3 Validation of approach: Soil moisture memory

The time series of simulated soil moisture produced with the optimal parameters, a reflection of precipitation, radiation and streamflow information only, is compared to the observed soil moisture in three highly monitored catchments to demonstrate that the precipitation, radiation, and streamflow data can indeed be translated into useful information on local soil moisture behavior. Because observed soil moisture information was not used at all in the calibration exercise, this comparison serves as a valid test of our methodology.

The validation focuses in particular on soil moisture persistence. There are many ways of quantifying soil moisture persistence; here, we compute it, for a given time of the year, as a lag correlation for a given lead (see *Koster and Suarez* 2001, *Seneviratne and Koster* 2012, and OS12) that ranges between 0 (no memory) and 1 (maximum memory). The memory we compute at a given day with a given time lag is defined as:

$$\rho(w_n, w_{n+t_{lag}}) = \frac{cov(w_n, w_{n+t_{lag}})}{\sigma_{w_n} \sigma_{w_{n+t_{lag}}}} \quad (8)$$

where $cov(w_n, w_{n+t_{lag}})$ denotes the covariance between soil moisture at days n and $n + t_{lag}$ in all considered years and σ_{w_n} refers to the standard deviation of soil moisture at day n using also the values of all considered years. Due to the limited available number of years of soil moisture observations (see Section 3.1), we in fact do some smoothing of the calculated persistences, computing representative estimates for half-monthly intervals. To determine the smoothed persistence for a given half-monthly interval, we use a "moving window" approach (OS12) that also considers the 30 days prior to the half-monthly interval and the 30 days after the end of the half-monthly interval. This can be expressed mathematically as:

$$\rho(w_n, w_{n+t_{lag}}) = \text{trimmed average} \left(\sum_{i=t_{start}-30}^{t_{end}+30-t_{lag}} \rho(w_i, w_{i+t_{lag}}) \right) \quad (9)$$

where t_{start} and t_{end} refer to the beginning and end of the particular half-monthly time period. The memory of that half monthly interval is then computed as a trimmed average of the $75 - t_{lag}$ individual persistences, avoiding days with the 10% largest and 10% lowest values (this last step differs from the approach of OS12, who take the median).

We compute the correlation for many different lags (from 1 to 40 days) in order to capture more completely the character of the persistence. One region may show high correlations at small lags and a rapid fall-off in correlation at longer lags, and another may show a fast fall-off at short lags and a slower fall-off thereafter; our computations will capture such differences in behavior. Thus, we calculate, from both observations and the simulations with optimized parameters, the correlation between soil moisture on a given date n with that at a later date $(n + t_{lag})$ across all years (Equation (9)). By computing a separate correlation for each date across all years, we avoid examining artificial memory associated with the climatological seasonal cycle of soil moisture. The higher the resulting correlation over a prescribed lag time, the higher we deem the soil moisture memory at that lag, and vice versa. Soil moisture memory is always decreasing with increasing time lag, because accumulated precipitation, runoff and ET alter the soil moisture content (Equation (1)). To facilitate the interpretation of soil moisture memory expressed as lag correlation, *Orth and Seneviratne* (2012b) compared the lag correlation with a persistence time scale (computed as mean duration to recover from anomalous conditions exceeding a certain threshold to normal conditions, expressed in days). They report an exponential relationship, i.e. the persistence time scale changes exponentially with linearly changing lag correlation.

Similarly to (9), but without time lag, we compute estimates for the standard deviations of e.g. initial soil moisture over all estimates of day n of all years. Using the moving window approach we obtain a number of estimates of which we take a trimmed average as a representative estimate for a particular half monthly interval.

Soil moisture persistences in this study are computed from April to October to exclude the

227 impact of snow cover, which is not included in the model. We therefore apply the model in an
228 extended period compared to the period July-September used for optimization to allow us to
229 show that the model also performs reasonably in months that are not used for calibration, but
230 that are still mostly snow-free (underlined by daily average temperatures above 0 degrees Celsius
231 on almost all days).

232 To compute uncertainties of the soil moisture memory estimates, we separate the whole time
233 period (24 years, see Section 3.2) into non-intersecting subsets of 3 years (period July-September
234 in each year as described in Section 2.2) and optimize the model in each catchment to yield one
235 parameter set per subset for a particular catchment. This is done with 5 repetitions for each
236 subset (instead of 20 used for the whole time series) due to computational constraints. We apply
237 all parameter sets of a particular catchment with the whole time series and derive respective
238 soil moisture memories; from these memories we then compute the standard deviations for every
239 considered month and lag time.

240 3 Data

241 3.1 Data analyzed for model validation

242 To validate the model, we use data from the three heavily monitored catchments: Oensingen
243 (Switzerland), Rietholzbach (Switzerland) and San Rossore (Italy). The climate at the Swiss sites
244 is temperate humid, whereas San Rossore is characterized by Mediterranean climate. Along with
245 the stream gauge measurements for the full catchments, there is a site in each catchment where
246 ET, radiation and precipitation have been recorded. Detailed information on the catchments and
247 sites is provided in Table 2.

248 ET at Rietholzbach was measured using a weighing lysimeter (*Seneviratne et al.* 2012), whereas
249 the eddy-covariance flux measurement method (*Baldocchi et al.* 2001) was used at the other two
250 sites. As this latter method is known for its energy-balance closure error (e.g. *Wilson et al.* 2002,

Foken et al. 2006, *Franssen et al.* 2010), we corrected the ET data with the following procedure: Using hourly values, we increased sensible and latent heat flux to equal net radiation while keeping the Bowen ratio constant (*Twine et al.* 2000). If the Bowen ratio was negative both fluxes were adjusted with respect to the strength of their dependence on net radiation instead. This strength was the slope obtained from the regression of all available values of the particular flux against net radiation on a particular time of the day; the flux with the higher slope was modified by the larger fraction of the energy balance deficit. Note that the ET data from San Rossore could not be corrected as no data of sensible heat flux is available for that site. Furthermore, we linearly detrended the soil moisture data from Rietholzbach to address a known problem with the sensors there (see also *Seneviratne et al.* 2012).

At all three catchments, we use satellite-derived net radiation data obtained from the NASA/GEWEX SRB project (http://eosweb.larc.nasa.gov/PRODOCS/srb/table_srb.html [checked on 26 March 2012]). Since these data only extend until 2007, we had to extrapolate net radiation from the available solar radiation measurements for the remaining 3 years at San Rossore. These were scaled to match the mean and standard deviation of the satellite net radiation of the previous 4 years. To evaluate the impact of this treatment we also applied such a scaling to solar radiation measured at Oensingen and found only minor impacts on the results there, predominantly on ET (not shown).

3.2 Data used for model application

Following validation, we apply the model to 13 near-natural catchments (i.e., catchments with little or no known human impact on streamflow) across Switzerland for which detailed stream gauge data are available. The catchments are located in a humid temperate climate, except for the Cassarate catchment in southern Switzerland where the climate is rather Mediterranean. A

summary of the catchment characteristics is provided in Table 3. The time period considered is 1984-2007.

For this period, we also obtained catchment-specific precipitation and radiation data. Precipitation forcing for the model was derived from several MeteoSwiss (Swiss Federal Office of Meteorology and Climatology) rain gauges in and/or near each respective catchment. The number of rain gauges per catchment depends on the size of the respective catchment and on the density of the network in the particular region (see <http://www.meteoschweiz.admin.ch/web/de/klima/messsysteme/boden.Par.0049.DownloadFile.tmp/karteniederschlagsmessnetz.pdf> [checked on 6 February 2013]). The measurements were weighted inversely according to their distance from the catchment in order to compute an area-representative estimate. As only solar radiation was measured at the ground, we used net radiation data from the NASA/GEWEX SRB project. A comparison of anomalies of the solar radiation measured at the ground with that from SRB showed correlations between 0.8 and 0.9 for the different catchments, underlining the good match also reported by OS12.

In order to study the dependency of soil moisture memory on topography (hilliness), we obtained values of mean compound topographic index (CTI; *Moore et al.* 1993) from the HYDRO-1K dataset [<http://webgis.wr.usgs.gov/globalgis/metadataqr/metadata/hydro1k.htm>]. As a measure of topography for each catchment, the CTI is a function of slope as well as upstream contributing area and increases with decreasing hilliness. Note that CTI is only evaluated at the catchments used for application of the model and not at the three validation catchments that include the very small Rietholzbach catchment. Therefore the 1km x 1km resolution is sufficient to characterize, to first order, the CTI amongst the catchments examined.

4 Results

In this section we first describe the application and validation of the simple model methodology in three heavily monitored catchments (Sections 4.2 and 4.3). We show its satisfactory ability to yield a realistic soil moisture memory despite its simplicity. In Section 4.4 we describe the application of the model in multiple catchments across Switzerland. This allows us to study the main meteorological controls of soil moisture memory as well as its dependency on altitude and topography.

4.1 Streamflow, runoff and evapotranspiration parameters

To summarize our methodology, we optimize the applied simple water balance model (i.e., we find optimized values for its five parameters) so that it reproduces well the time variations in the daily streamflows measured in a given catchment when forced with local precipitation and net radiation data. An overview of the fitted parameters in all catchments is provided in Table 4.

A note about the parameter search is appropriate here. Two parameters, the ET ratio exponent and maximum ET ratio, collide with their bounds in 2 and 9 catchments, respectively, out of the 16 catchments considered in total in this study (see Table 1 for bounds and Table 4 for fitted parameters). Concerning the maximum ET ratio, the fact that the optimum value of β_0 is found to be exactly 1, an imposed bound for the parameter, does not reflect poorly on the parameter estimation approach; the optimized value of 1 simply means that for the catchment in question, all of the net radiation is converted to evaporation in wet conditions, a physically plausible scenario. Our requirement that β_0 cannot exceed 1 is simply a reflection of our assumption that net radiation provides the energy needed for evaporation. Whereas high maximum ET ratios are not surprising in a radiation-limited regime that is characteristic for Switzerland, we note the possibility that in nature, ET might (temporarily) exceed net radiation through processes that

are not captured by our simple model, such as energy input from warm air advection to Europe; therefore, the collisions experienced with the β_0 term can be said to reflect the limitations of our assumption. Note that both parameters that collide with their bounds are related to radiation (Equation 2) and streamflow (through optimization procedure, see Section 2.2). Therefore it is furthermore possible that these collisions are due to scale discrepancies between radiation and streamflow measurements and the consequent mismatch in their temporal evolutions. However, the parameter collisions should, in any case, not have a major impact on the resulting estimated soil moisture memory as indicated by supplemental tests (not shown) in which the bounds were removed (in conflict with the model's underlying assumptions) and the results were found to be generally similar. Furthermore we note that despite the parameter collisions we find a good match between modeled and observed soil moisture memory as described in the following subsection.

To validate our optimization procedure, we applied it with higher (coarser) step widths for the parameters and then compared the results with those obtained when all possible combinations of parameters (assuming the same coarse spacing) were tested. This allowed us to compare the resulting best parameter sets. Given the high computational effort the validation was done only for the three catchments listed in Table 2. The best parameter sets found from both procedures were identical for all three catchments (see Table 4 for parameter values), underlining the validity of the approach introduced in this study. As expected due to the larger step width (lower accuracy), these parameter sets yield slightly lower correlations between observed and modeled streamflow compared to the parameter sets found using the default step widths (see Table 1).

4.2 Validation of estimated memory

In addition to generating realistic streamflows, the optimized model produces, as a matter of course, a time series of daily soil moisture, from which soil moisture persistence measures can be derived. Our methodology for converting streamflow measurements into soil moisture information

is deemed successful if the derived soil moisture persistences obtained from this time series agree with those obtained using independent soil moisture measurements in the catchments.

This validation test was performed in each of the three catchments described in Section 3.1. Results are shown in Figure 1. Shown for each catchment are the modeled and observed persistences for different lags (out to 40 days) and for different times of the year (April through October). Overall, the memory characteristics in the three catchments are well captured by the model, with a reasonable representation in each of the seasonal cycle of soil moisture memory and its decay with lag. The observed and simulated memory is comparatively strong at Oensingen and San Rossore and weakest at Rietholzbach. The seasonal cycle of the observed memory at San Rossore differs clearly from that of the other two sites, and this is captured by the model. Difference plots are shown in the bottom row of the figure; there is no clear pattern of over- and underestimation of memory in the simulation results. The relatively large difference between modeled and observed soil moisture memory in autumn at both Rietholzbach and San Rossore is consistent with results of OS12 (Figure 6 of that paper), who used the same atmospheric forcing data, and derived also a clearly underestimated soil moisture memory. In this previous study, the identified reason for this behavior was a mismatch between precipitation and soil moisture observations in autumn at these two sites. This means that the water balance is not closed with the employed observations, which could be due, for example, to a higher spatial variability of precipitation or a stronger role of land cover in this season.

The model, using only information on locally measured precipitation, net radiation, and streamflow, therefore successfully captures the distinctions between the catchments in their soil moisture memory behavior. Despite its simplicity, it captures enough of the physical processes controlling memory to allow the translation of streamflow information into soil moisture information. Furthermore, the agreement in Figure 1 suggests (as does the reasonable reproduction of soil moisture anomalies shown below) that the time behavior of the observed site-based soil moisture

anomalies is representative of that for soil moisture across the catchment containing the site; that is, soil moisture levels may be spatially heterogeneous within a catchment but may nevertheless show similar temporal dynamics. This is consistent with results from *Mittelbach and Seneviratne* (2012) for Switzerland based on measurements from the Swiss Soil Moisture Experiment (SwissS-MEX), which show that soil moisture dynamics have a large regional footprint in that region, unlike absolute soil moisture that displays a stronger spatial variability.

To illustrate further the impact of the fitted parameter set on the resulting soil moisture memory characteristics – in particular, to show the relative impacts on memory of the parameter values and the meteorological forcing – we run the model at each of the three sites mentioned above with the parameter set fitted for the particular site and also with the parameter sets fitted for the other two sites. The results are displayed in Figure 2. We find that the parameter set is more important for determining the resulting soil moisture memory than is the meteorological forcing. There are similarities between the actually modeled memory at Oensingen and San Rossore and the resulting memory when using the parameter set or meteorological forcing from another site. This can be explained by the roughly similar fitted parameters (see Table 4). Generally the strong sensitivity of the memory with respect to the parameter set underlines the ability of our simple model framework to yield a parameter set that is related with realistic features of the studied catchments.

4.3 Hydrological states and fluxes

While the main goal of the tested methodology is the extraction of soil moisture memory statistics, we can also validate the soil moisture, streamflow, and evapotranspiration time series produced by the optimized model against available observations in the three validation catchments. Comparisons of the observed and simulated anomalies of these quantities are provided in the top three rows of Figure 3. Mean seasonal cycles have been subtracted from both the observed and

simulated data in order to avoid an overestimation of model skill associated with the seasonal cycles inherent in the precipitation and net radiation forcing. While this subtraction also prevents a proper evaluation of bias, such bias evaluations would, in any case, be of limited usefulness: (i) significant biases are likely in the observed evaporation data given the closure problem associated with eddy covariance measurements (see Section 3.1), (ii) biases in soil moisture are likely because the model uses an arbitrary wilting point (which doesn't affect the temporal variability of the soil moisture it produces), (iii) observed absolute soil moisture is also expected to vary strongly even on small spatial scales, only the temporal dynamics should display a regional footprint (*Mittelbach and Seneviratne* 2012), and (iv) biases in streamflow may occur especially in flat catchments through baseflow out of the catchment away from the stream gauge.

Overall, the model seems to do especially well in estimating soil moisture variations, particularly for Oensingen ($R^2 = 0.78$) but also for Rietholzbach ($R^2 = 0.62$). Streamflows for these two sites are also reasonably reproduced (R^2 values of 0.6 and 0.87, respectively), whereas simulated ET values are somewhat less consistent with the observations, although still satisfactory at Rietholzbach ($R^2 = 0.58$). The simulated values are always worse for the San Rossore catchment, possibly due to (i) its larger size and the corresponding reduction in the large-scale representativeness of its site-based precipitation forcing and (ii) the interpolation of the radiative forcing (see lower part of Section 3.1). In cases of comparatively low R^2 values, such as for ET at Oensingen and for all quantities at San Rossore, we find that the model tends to underestimate the variability of the anomalies, as indicated by the regression slopes that are clearly smaller than 1.

Corresponding scatter plots produced with data from June and October (not shown) show comparable agreement between the model results and observations. This provides an independent evaluation of model performance, given that these months were not part of the fitting period (see Section 2.2).

The bottom row in Figure 3 displays the optimized runoff functions (solid red lines) and ET

functions (solid black lines) at Oensingen, Rietholzbach and San Rossore. Every plotted point represents either an observed streamflow ratio, $\frac{S_n}{P_n}$ (in red), or an observed ET ratio, $\frac{\lambda \rho_w E_n}{R_n}$ (in black), with the respective quantities (both the numerators and the denominators separately) accumulated over a week to increase representativeness and to ensure comparability between runoff ratio as shown by the fitted function and streamflow ratio as shown by the observations.

At first glance, the evaporation functions seem to disagree with the data. Here one must remember two key points: (i) no evaporation data were used in the optimization of the functions, and (ii) the evaporation observations are subject to bias and, even after bias correction, are uncertain. Eddy-covariance measurements are known, for example, to produce underestimated fluxes (e.g. *Wilson et al.* 2002, *Foken et al.* 2006, *Franssen et al.* 2010). Therefore we corrected the ET in order to close the energy balance through a modification of latent and sensible heat fluxes as described in Section 3.1. Indeed, at Rietholzbach, where ET was measured with a weighing lysimeter instead, the modeled ET ratio compares better to observations. At San Rossore, the ET data could not be corrected because net radiation was not available over the whole time period; the observed ET fluxes there are thus underestimated. (Such errors might also explain the relatively poor comparison of ET anomalies at San Rossore in the third row of Figure 3.) At Oensingen, ET flux corrections may have led to excessive ratios, possibly because ET was measured over grassland whereas the optimized function represents the whole catchment, which includes forested regions. *Teuling et al.* (2010b), using observations, showed that forests in temperate Europe use water more conservatively than grassland, especially under extreme conditions (Figure 1 of that paper).

In contrast, the optimized runoff functions do capture, to first order, the observed streamflow ratios. This makes sense, given that the streamflow measurements were used in the optimization procedure. The high fitted runoff ratio (especially for wet conditions) corresponds well with the generally wet regime at Rietholzbach (annual precipitation $\approx 1500\text{mm}$), such that most of the precipitation can not be stored but runs off instead. There is nevertheless still some bias seen in the optimal runoff functions and a substantial amount of scatter seen in the streamflow

ratio observations. Again, our use of available streamflow and precipitation observations is made difficult by the mismatch in their scales; because the functions are optimized using data from July to September, we speculate that local thunderstorms and showers might influence parts of a catchment not captured by the rain gauge or might over-emphasize small-scale storms falling over the rain gauges. Of course, even without a scale mismatch, scatter in the plotted points will result from the fact that precipitation and streamflow measurements each have their own errors, and these errors are compounded when the ratio is computed. Again, some time shift between precipitation and streamflow is already implicitly included in Equations 4-6 through the streamflow recession.

Summing up we note that generally the unimpressive agreement found in the bottom row in Figure 3 is no surprise given the vastly different scales we consider (e.g. for streamflow and precipitation or of modeled, catchment-scale ET and observed, point-scale ET) and the noted measurement uncertainties related to, for example, eddy-covariance ET measurements or point-scale precipitation measurements. When considering this unimpressive agreement, it is worth remembering that the optimization procedure focuses on finding the runoff and evaporation functions that best reproduce the time variability of the observed streamflow (through an R^2 value), a reflection of the time dynamics of the local hydrological cycle, rather than functions that are necessarily consistent with direct evaporation and streamflow measurements, as represented by the plotted points in the lowest row of Figure 3. Naturally, if the latter approach were used, the functions chosen would agree much more strongly with those plotted points. Of course, the latter approach requires soil moisture and evaporation information, which is what we want to avoid here, given the noted dearth of contemporaneous soil moisture and evaporation data. While it is certainly possible that our optimization approach does not produce the runoff function and ET function combination that best reproduces the measured soil moisture memory, it does nevertheless produce a combination that reproduces it reasonably well (Figure 1), and it does maintain the critical advantage of being based on only streamflow, precipitation, and radiation information.

Note furthermore that the suitability of the optimization approach may vary depending on the climate regime, as it becomes difficult, under dry conditions when streamflow variations are small, to infer hydrological variability of a catchment from streamflow only (e.g. *Teuling et al.* 2010a).

For completeness, Figure 4 shows the hydrographs associated with the optimized values of τ for the three catchments. In the Rietholzbach catchment, the streamflow response falls off most quickly, as might be expected given the catchment's hilliness and relatively small size. In the other two catchments, 2% of the water in a precipitation event is still running off two weeks after the event.

4.4 Application to multiple Swiss catchments

The application of the methodology to precipitation, net radiation, and streamflow data in 13 catchments across Switzerland (Section 3.2) allows us to obtain an areal picture of soil moisture memory (30-day-lagged autocorrelation), as shown in Figure 5. The similar memories found for adjacent catchments, even those with different sizes, provides additional support for our approach. A signature of the alpine ridge (and its associated precipitation regime) is seen in the memory distribution.

The highest memory is found for the Langeten catchment, which is located in the Swiss plateau between the Alps and the Jura mountains. High memory is also found for the Mentue (also in the Swiss plateau) and Ergolz (northern end of the Jura mountains) catchments. The lowest memory is found in the highest catchments: Sitter, kleine Emme, Emme and Sense. Overall, soil moisture memory seems to increase with increasing distance from the Alps, as seen in the far west for the Broye and Mentue catchments and in the far east for the Murg, Aach, Goldach and Sitter catchments. Despite the drier climate regime south of the Alps, we find a similar strength of the soil moisture memory at Cassarate compared to catchments along the northern alpine front.

Figures 6 and 7 summarize the results for all catchments, showing the optimized runoff and ET

functions (Column 1) and the corresponding soil moisture memories as a function of season and lag (Column 2). The rows holding the catchment results are arranged in order of average memory, starting with Langeten (the catchment with the strongest memory). The optimized functions differ significantly among the catchments, as does the absolute soil moisture range. Correspondingly, the strength of the estimated soil moisture memory and its seasonal cycle differ significantly across the catchments, especially in summer. In general, memory seems to be strongest in autumn, for which considerable memory is often seen at 4-5 week lags, and it is weakest in spring, which generally shows almost no significant memory beyond 2 weeks.

Figures 6 and 7 also display the uncertainties corresponding to the soil moisture memories, as derived with the methodology described in Section 2.3. They are mostly smaller than 0.2, indicating that the computed memory patterns are robust with respect to parameter sets obtained from different and independent subsets of the full time period analyzed. Especially if the estimated memory is high, the uncertainties are low; therefore high soil moisture memory as identified with the simple water balance model is particularly reliable.

4.4.1 Controls of soil moisture memory in Switzerland

OS12 identified two main controls of soil moisture memory at five sites in Central and Mediterranean Europe; (i) the ratio between the variability of initial soil moisture and subsequent forcing, and (ii) the correlation between initial soil moisture and the subsequent forcing. They also report that the forcing is dominated by precipitation, and thus we can express the first control as the unitless ratio between the standard deviation of initial soil moisture and the standard deviation of subsequent precipitation:

$$\tilde{\kappa}_n = \frac{\sigma_{w_{n,y}}}{\sigma_{P_{n,y}t_{lag}}} \quad (10)$$

where $P_{n,y}$ denotes precipitation (in $\frac{mm}{d}$) between date n and $n + t_{lag}$ of year y . It is multiplied with t_{lag} to yield the accumulated precipitation during that interval. The standard deviations are

computed as described in Section 2.3. Note that $\tilde{\kappa}_n$ also reflects the impact of seasonal variations in precipitation. Given that precipitation dominates the forcing, the second control identified in OS12 can be simplified to yield $\rho(w_n, P_n)$. High values of either of these controls are indicative of higher soil moisture memory. The first control, $\tilde{\kappa}_n$, reflects the size of the anomaly to be erased relative to that of the precipitation available to erase it, and the second describes how the effect of the precipitation may be diminished if its magnitude is not independent of the initial anomaly.

The fourth and fifth columns of Figures 6 and 7 illustrate the values of these controls at all catchments, for all months and lags considered. The ratio of the soil moisture and precipitation variabilities decreases from the top to the bottom in both figures as the soil moisture memory decreases, suggesting a connection. Confirming the results of OS12, also comparatively high correlations between initial soil moisture and the subsequent precipitation (a reflection, indeed, of memory in precipitation itself) seem to coincide with high memory in most catchments.

Moreover these two figures show that the runoff optimization approach (Section 2.2) yields functions of similar shape for nearby catchments (e.g. Mentue/Broye and Sitter/Goldach), underlining the robustness of the simple model approach. However, despite such similarity in the functions, we can sometimes find different strengths for the soil moisture memory, as in the Mentue and Broye catchments, illustrating the importance of catchment-specific parameters such as water holding capacity and maximum ET ratio.

Figure 8 shows the correlations between the optimized model parameters and the resulting soil moisture memory (as shown in Figure 5). Water holding capacity is seen to be a strong control of soil moisture memory, which is intuitively sensible; it has a direct impact on the numerator of the standard deviation ratio discussed above. A second control of memory is the runoff ratio exponent (even if of questionable statistical significance due to the relatively small set of catchments). The higher this exponent, the greater the contrast in the impact of runoff on soil moisture in wet and dry conditions. That is, for a high exponent, the dampening impact of runoff on soil moisture anomalies is significantly reduced in drier conditions. OS12 found that especially dry anomalies

contribute to a higher soil moisture memory, which explains why the runoff ratio exponent has such a large effect. It is important to note that these three controls are not statistically independent, for otherwise they would explain too much of the variance of soil moisture memory; a proper breakdown of the roles of these parameters and how they vary with each other would require a substantially larger collection of analyzed catchments.

4.4.2 Dependence of soil moisture memory on altitude, topography and dryness index

Investigating the dependency of soil moisture memory on altitude, topography and dryness index allows us to separate the effects of soil and vegetation characteristics, morphology and atmospheric forcing, respectively. Even if altitude and topography are usually related, here they are to some extent independent due to the complex (pre-) alpine terrain of Switzerland. The top row of Figure 9 demonstrates that memory decreases with altitude and with increased topography (expressed as CTI, see Section 3.2). The bottom row of the figure shows a link between $\tilde{\kappa}_n$ (the aforementioned ratio of initial soil moisture variability to precipitation variability shown in Equation (10)) and both altitude and topography. This $\tilde{\kappa}_n$ ratio was identified in Section 4.4.1 as a main control of soil moisture memory, which is consistent with the shown dependencies on altitude and topography. The higher (or hillier) a catchment is, the thinner the soil should be, leading to a decreased water holding capacity and therefore a lower σ_{w_n} and thus a lower $\tilde{\kappa}_n$ value. Even if topography and altitude are found to have the same impact on soil moisture memory, the reasons may not be the same, since topography as such only impacts soil moisture dynamics whereas altitude also reflects the varying atmospheric forcing (e.g. precipitation (variability) increasing with altitude and thereby reducing soil moisture memory as described in the previous subsection).

We also investigated the link between mean soil moisture memory (as shown in Figure 5) and catchment-specific dryness index, as illustrated with the plots on the right hand side of Figure 9. The dryness index is computed as $\frac{\bar{R}}{\lambda \rho_w \bar{P}}$, where λ is the latent heat of vaporization and \bar{R} and

\bar{P} are long-term averages of annual net radiation and precipitation, respectively. Soil moisture memory tends to increase with increasing dryness index, even if the diagnosed relationship between the two is rather weak. Less precipitation leads to a lower variability and thus a higher $\tilde{\kappa}_n$ value as shown in the Figure.

Comparing the influence of these three controls on soil moisture memory in Switzerland as indicated by the R^2 values, we find that altitude is of highest importance, followed by topography and dryness index.

5 Conclusions

In this study we modified the simple water-balance model proposed by *Koster and Mahanama* (2012) to include such features as streamflow recession and an implicit form of the water balance equation. We then applied the model to the analysis of soil moisture memory. Our main tested hypothesis was whether such a simple model can be used to extract information on soil moisture memory based on observations of precipitation, net radiation, and streamflow alone, since these observations are much more plentiful than soil moisture observations.

Our approach was successfully validated using data from some of the relatively rare catchments for which soil moisture measurements and contemporaneous meteorological measurements are adequate. Using only precipitation, net radiation, and streamflow data, the model captures the first order behavior of the observed soil moisture memory in terms of its variation with season and the considered lag (Figure 1). The model also reproduces the observed soil moisture anomalies reasonably well (Figure 3).

We then used the validated model to estimate the soil moisture memory within 13 near-natural catchments across Switzerland. The resulting spatial distribution of estimated memory allowed an analysis of the controls on this memory. Our results support earlier propositions that the main controls of memory in Central Europe are (i) the ratio of the standard deviations of initial soil

moisture and subsequent precipitation, and (ii) the correlation between the initial soil moisture and the subsequent precipitation. Soil moisture memory in the vicinity of the Alps appears to decrease with altitude and hilliness (as measured by CTI), possibly because soils at higher elevations tend to be thinner.

The study with the 13 Swiss catchments demonstrates that the simple water balance model can be used in conjunction with precipitation, net radiation, and streamflow measurements to estimate soil moisture memory and its controls even in the absence of direct soil moisture measurements. Applying this methodology to catchments in other regions of the world could help identify areas of strong soil moisture memory, that is, areas for which soil moisture initialization has a chance to contribute to hydrological or meteorological prediction.

Acknowledgments

We acknowledge the Swiss Federal Office for the Environment (FOEN) for providing streamflow data for the 13 Swiss catchments the the Swiss weather service (MeteoSwiss) for providing the corresponding precipitation data. We thank Massimiliano Zappa for sharing these data with us, as well as Guenther Seufert and Christoph Ammann for providing data from San Rossore and Oensingen, respectively. We also thank three anonymous reviewers for helpful comments on the manuscript and Gerd Vogel and Guenther Seufert for advice on the CEOP and Carboeurope databases, which we acknowledge for sharing the Falkenberg, Kehrigk and San Rossore data, as well as the hydrological service of the Tuscany Region for providing the streamflow data of the Arno river. We acknowledge financial support from the Swiss National Foundation through the NRP61 DROUGHT-CH project, as well as partial support from the EU-FP7 DROUGHT-RSPI project.

Appendix A: Optimization procedure for identification of catchment-specific parameter sets

We first choose a random value for each parameter in Equations (2) and (6) from within a prescribed acceptable range and add a prescribed step width (see Table 1) to yield a second value for each parameter. We then run the model for all $2^5 = 32$ combinations of parameters to find the set which yields the highest correlation between modeled and observed streamflow. After that, we rerun the model using another $2^5 = 32$ combinations, assigning to each parameter the optimal value found before and this value with the respective step width subtracted (if the lower value from before was the optimal value) or added (if the higher value from before was the optimal value). This procedure is repeated until the same set of parameters is found two times in a row.

This procedure, of course, guarantees only a local (rather than a global) optimum in the five-dimensional parameter space. We thus repeat the procedure 20 times, always starting with new randomly chosen values for each parameter. This yields 20 local optima, of which many are similar or even identical, underlining the robustness of this approach. Of these 20 local optima we take the best as our parameter set for a given catchment. Our tests with the procedure suggest that higher computational effort would probably not yield a different solution; given the step widths applied to the parameters, we most likely indeed find the global optimum in the five-dimensional parameter space.

634 References

- 635 Ammann, C., O. Marx, V. Wolff, and A. Neftel (2010), Measuring the biosphere-atmosphere ex-
 636 change of total reactive nitrogen by eddy covariance using a novel converter, in *29th Conference*
 637 *on Agricultural and Forest Meteorology*.
- 638 Baldocchi, D. D., et al. (2001), FLUXNET: A new tool to study the temporal and spatial variability
 639 of ecosystem-scale carbon dioxide, water vapor and energy flux densities, *Bull. Amer. Meteorol.*
 640 *Soc.*, 82, 2415–2435.
- 641 Dorigo, W. A., et al. (2011), The international soil moisture network: a data hosting facility for
 642 global in situ soil moisture measurements, *Hydrol. Earth Syst. Sci.*, 15, 1675–1698.
- 643 Douville, H. (2010), Relative contribution of soil moisture and snow mass to seasonal climate
 644 predictability: A pilot study., *Clim. Dyn.*, 34, 797–818.
- 645 Entin, J. K., A. Robock, K. Y. Vinnikov, S. E. Hollinger, S. Liu, and A. Namkhai (2000), Temporal
 646 and spatial scales of observed soil moisture variations in the extratropics, *J. Geophys. Res.*, 105,
 647 11,865–11,877.
- 648 Foken, T., F. Wimmer, M. Mauder, C. Thomas, and C. Liebethal (2006), Some aspects of the
 649 energy balance closure problem, *Atmos. Chem. Phys.*, 6, 4395–4402.
- 650 Franssen, H. H., R. Stöckli, I. Lehner, E. Rotenberg, and S. I. Seneviratne (2010), Energy balance
 651 closure of eddy covariance data: a multi-site analysis for European FLUXNET stations, *Agric.*
 652 *Forest Meteorol.*, 150 (12), 1553–1567.
- 653 Hirschi, M., S. I. Seneviratne, V. Alexandrov, F. Boberg, C. Boroneant, O. B. Christensen, H. For-
 654 mayer, B. Orlowsky, and P. Stepanek (2011), Observational evidence for soil-moisture impact
 655 on hot extremes in southeastern Europe, *Nature Geoscience*, 4, 17–21.
- 656 Kirchner, J. (2009), Catchments as simple dynamical systems: Catchment characterization,
 657 rainfall-runoff modeling, and doing hydrology backward, *Water Resources Research*, 45,
 658 W02,429.
- 659 Koster, R. D., and S. Mahanama (2012), Land Surface Controls on Hydroclimatic Means and
 660 Variability, *J. Hydrometeorol.*, 13, 1604–1620.
- 661 Koster, R. D., and M. J. Suarez (2001), Soil moisture memory in climate models, *J. Hydrometeo-*
 662 *rol.*, 2, 558–570.
- 663 Koster, R. D., S. P. P. Mahanama, B. Livneh, D. P. Lettenmaier, and R. H. Reichle (2010a), Skill
 664 in streamflow forecasts derived from large-scale estimates of soil moisture and snow, *Nature*
 665 *Geoscience*, 3, 613–616.

- 666 Koster, R. D., et al. (2004), Regions of strong coupling between soil moisture and precipitation,
667 *Science*, *305*, 1138–1140.
- 668 Koster, R. D., et al. (2010b), Contribution of land surface initialization to subseasonal forecast
669 skill: First results from a multi-model experiment, *Geophys. Res. Lett.*, *37*, L02402.
- 670 Mahanama, S. P. P., B. Livneh, R. D. Koster, D. Lettenmaier, and R. Reichle (2012), Soil
671 moisture, snow, and seasonal streamflow forecasts in the United States, *J. Hydrometeorol.*, *13*,
672 189–203.
- 673 Mittelbach, H., and S. I. Seneviratne (2012), A new perspective on the spatio-temporal variability
674 of soil moisture: temporal dynamics versus time invariant contributions, *Hydrol. Earth Syst.*
675 *Sci.*, *16*, 2169–2179.
- 676 Moore, I. D., P. E. Gessler, G. A. Nielsen, and G. A. Petersen (1993), Terrain attributes: estimation
677 methods and scale effects, in: *Modeling Change in Environmental Systems*, pp. 189–214.
- 678 Mueller, B., and S. I. Seneviratne (2012), Hot days induced by precipitation deficits at the global
679 scale, *Proc. Natl. Acad. Sci.*, *109* (31), 12,398–12,403.
- 680 Orth, R., and S. I. Seneviratne (2012a), Analysis of soil moisture memory from observations in
681 Europe, *J. Geophys. Res.*, *117*, D15,115, doi:10.1029/2011JD017366.
- 682 Orth, R., and S. I. Seneviratne (2012b), Propagation of soil moisture memory to runoff and
683 evapotranspiration, *Hydrol. Earth Syst. Sci. Disc.*, *9*, 12,103–12,143.
- 684 Robock, A., K. Y. Vinnikov, G. Srinivasan, J. K. Entin, S. E. Hollinger, N. A. Speranskaya, S. Liu,
685 and A. Namkhai (2000), The global soil moisture data bank, *Bull. Amer. Meteorol. Soc.*, *81*,
686 1281–1299.
- 687 Seneviratne, S. I., and R. D. Koster (2012), A revised framework for analyzing soil moisture
688 memory in climate data: Derivation and interpretation, *J. Hydrometeorol.*, *13*, 404–412.
- 689 Seneviratne, S. I., T. Corti, E. L. Davin, M. Hirschi, E. B. Jaeger, I. Lehner, B. Orlowsky, and
690 A. J. Teuling (2010), Investigating soil moisture-climate interactions in a changing climate: A
691 review, *Earth-Science Reviews*, *99*, 125–161.
- 692 Seneviratne, S. I., et al. (2006), Soil moisture memory in AGCM simulations: Analysis of Global
693 Land-Atmosphere Coupling Experiment (GLACE) data, *J. Hydrometeorol.*, *7*, 1090–1112.
- 694 Seneviratne, S. I., et al. (2012), The Rietholzbach research site: Analysis of 32-year hydrocli-
695 matological time series and 2003 drought at a Swiss pre-alpine catchment, *Water Resources*
696 *Research*, *48*, W06,526.
- 697 Teuling, A. J., I. Lehner, J. W. Kirchner, and S. I. Seneviratne (2010a), Catchments as simple
698 dynamical systems: experience from a Swiss pre-alpine catchment, *Water Resources Research*,
699 *46*, W10,502.

- 700 Teuling, A. J., et al. (2010b), Contrasting response of European forest and grassland energy
701 exchange to heatwaves, *Nature Geoscience*, 3, 722–727.
- 702 Tirone, G. (2003), Stima del bilancio del carbonio di due ecosistemi forestali Mediterranei. Con-
703 fronto tra una lecceta e una pineta., Ph.D. thesis, University of Tuscia, Viterbo, Italy. [in Italian].
- 704 Twine, T. E., W. P. Kustas, J. M. Norman, D. R. Cook, P. R. Houser, T. P. Meyers, J. H. Prueger,
705 P. J. Starks, and M. L. Wesely (2000), Correcting eddy-covariance flux underestimates over a
706 grassland, *Agric. For. Meteorol.*, 103, 279–300.
- 707 Vinnikov, K. Y., and I. B. Yeserkepova (1990), Soil moisture: Empirical data and model results,
708 *J. Climate*, 4, 66–79.
- 709 Viterbo, P., and A. K. Betts (1999), Impact of the ECMWF reanalysis soil water on forecasts of
710 the July 1993 Mississippi flood, *J. Geophys. Res.*, 19, 361–366.
- 711 Wilson, K., et al. (2002), Energy balance closure at FLUXNET sites, *Agric. Forest Meteor.*, 113,
712 223–243.

Table 1: Overview of step width of model parameters as used in the optimization procedure, of their boundaries and the range of their respective estimates.

Parameter	Step width	Lower limit	Upper limit	Maximum value found	Minimum value found
water holding capacity c_s (mm)	30	30	-	500	80
inverse streamflow recession timescale $\frac{1}{\tau}$ (1/days)	0.02	0.02	-	0.80	0.10
runoff ratio exponent α	0.2	0.2	-	8.0	0.8
ET ratio exponent γ	0.03	0.03	-	1.05	0.03
max ET ratio β_0	0.03	0.03	0.99	0.99	0.60

Table 2: Overview of measurements and conditions at the sites and catchments used for validating the model as well as references describing the sites in more detail.

Station	Data period	Land cover	Soil type	SM measurement depths (m)
Oensingen (CH)	2002 - 2007	grassland	clay	0.05, 0.1, 0.3, 0.5
Rietholzbach (CH)	1994 - 2007	grassland	(clay) loam	0.05, 0.15, 0.55
San Rossore (ITA)	2004 - 2010	forest	sand	0.1, 0.3, 0.45

Station	Streamflow station (distance and direction rel. to SM station in km)	Catchment area	Satellite radiation coordinates	Reference
Oensingen (CH)	Brugg (38 east)	11726km ²	47.5°N 7°E	<i>Ammann et al.</i> 2010
Rietholzbach (CH)	Mosnang (1.5)	3.3km ²	47.5°N 9°E	<i>Seneviratne et al.</i> 2012
San Rossore (ITA)	Vicopisano (25 east)	8228km ²	43.5°N 11°E	<i>Tirone</i> 2003

Table 3: Overview of catchments where the model is applied.

Catchment	Size (km ²)	Mean altitude (m above sea level)	Mean CTI	Mean daily stream-flow (mm)	Satellite radiation coordinates
Aach	49	480	11.82	1.32	47.5°N 9°E
Broye	392	710	11.33	1.78	46.5°N 7°E
Cassarate	74	990	9.39	2.72	45.5°N 9°E
Emme	124	1189	10.03	3.01	46.5°N 7°E
Ergolz	261	590	10.99	1.25	47.5°N 7°E
Goldach	50	833	10.71	2.32	47.5°N 9°E
Guerbe	54	837	9.98	2.01	46.5°N 7°E
kleine Emme	477	1050	10.48	2.81	46.5°N 7°E
Langeten	60	766	11.37	1.79	47.5°N 7°E
Mentue	105	679	11.27	1.34	46.5°N 7°E
Murg	79	650	11.47	1.98	47.5°N 9°E
Sense	352	1068	10.5	2.18	46.5°N 7°E
Sitter	74	1252	10.18	4.06	47.5°N 9°E

Table 4: Overview of fitted parameters for all catchments.

	Catchment	water holding capacity c_s (mm)	inverse streamflow recession timescale $\frac{1}{\tau}$ (1/days)	runoff ratio exponent α	ET ratio exponent γ	max ET ratio β_0
Catchments in which model is validated	Oensingen	410	0.10	0.8	0.03	0.60
	Rietholzbach	140	0.80	4.4	0.42	0.99
	San Rossore	500	0.14	3.6	0.03	0.96
Catchments in which model is applied	Aach	230	0.62	8.0	0.78	0.99
	Broye	200	0.36	5.8	0.42	0.60
	Cassarate	410	0.36	6.8	0.33	0.81
	Emme	80	0.74	1.4	0.27	0.99
	Ergolz	290	0.54	5.6	0.90	0.99
	Goldach	350	0.60	6.8	0.75	0.99
	Guerbe	170	0.44	4.2	1.05	0.99
	kleine Emme	80	0.66	2.4	0.60	0.99
	Langten	320	0.52	4.0	0.06	0.81
	Mentue	410	0.52	6.4	0.66	0.99
	Murg	230	0.50	6.2	0.63	0.99
	Sense	80	0.52	1.6	0.09	0.69
	Sitter	170	0.56	7.4	0.90	0.69

Figure 1: Soil moisture memory computed from observed and modeled soil moisture in the 3 validation catchments for lag times between 5 and 40 days. Values outside the plotting range of the difference plots are shaded in gray.

Figure 2: Soil moisture memory computed for all possible combinations of meteorological forcings and parameter sets from the 3 validation catchments as compared to observed soil moisture memory displayed in the bottom row.

Figure 3: The top rows show modeled soil moisture, streamflow and evapotranspiration plotted against observations for data within the period July-September that was used to fit the functions. The red lines are fitted through least-squares regressions. The bottom row shows the functions of Equations (2) (black) and (6) (red) fitted for each catchment. These are compared to weekly-averaged observed corresponding ratios plotted as points against observed soil moisture (mean and variance adapted to model soil moisture).

Figure 4: Fitted hydrographs (Equation (4)) in the 3 validation catchments.

Figure 5: Soil moisture memory of lag 30 days at all investigated catchments across Switzerland, averaged from April through October. The brownish background indicates the topography, with darker brown referring to higher altitudes.

Figure 6: Overview of fitted functions, soil moisture memory, its uncertainty (refer to text for details), $\tilde{\kappa}_n = \frac{\sigma_{w_n, y}}{\sigma_{P_n, y^{t_{lag}}}}$ and $\rho(w_n, P_n)$ (as described in Section 4.4.1) for all catchments going from high soil moisture memory (top row) to low soil moisture memory (bottom row). In the left hand side column the red curves correspond to the fitted runoff ratio functions, the black lines show the fitted ET ratio functions and the vertical blue lines denote the 5% and 95% quantile of

all soil moisture values in the time frame between April and October. Soil moisture memory, its uncertainty, $\tilde{\kappa}_n$ and $\rho(w_n, P_n)$ are computed for all months between April and October and for lag times between 5 and 40 days.

Figure 7: Continuation of Figure 6.

Figure 8: Correlations of fitted model parameters (listed in Table 1) at all catchments with respective soil moisture memory at a lag of 30 days. Dark gray corresponds to negative correlations, light gray indicates positive correlations. Hatching indicates correlations that are not significant on the 5% level (two-sided t-test).

Figure 9: The top row displays the soil moisture memories of lag 30 days of all 13 catchments plotted against altitude, CTI and dryness index including a least-squares fit and explained fraction of variance. The same is shown in the lower row for the ratio $\tilde{\kappa}_n = \frac{\sigma_{w_n, y}}{\sigma_{P_n, y} t_{lag}}$, also for a lag time of 30 days.




Figure 1: Soil moisture memory computed from observed and modeled soil moisture in the 3 validation catchments for lag times between 5 and 40 days. Values outside the plotting range of the difference plots are shaded in gray.




Figure 2: Soil moisture memory computed for all possible combinations of meteorological forcings and parameter sets from the 3 validation catchments as compared to observed soil moisture memory displayed in the bottom row.




Figure 3: The top rows show modeled soil moisture, streamflow and evapotranspiration plotted against observations for data within the period July-September that was used to fit the functions. The red lines are fitted through least-squares regressions. The bottom row shows the functions of Equations (2) (black) and (6) (red) fitted for each catchment. These are compared to weekly-averaged observed corresponding ratios plotted as points against observed soil moisture (mean and variance adapted to model soil moisture).




Figure 4: Fitted hydrographs (Equation (4)) in the 3 validation catchments.




Figure 5: Soil moisture memory of lag 30 days at all investigated catchments across Switzerland, averaged from April through October. The brownish background indicates the topography, with darker brown referring to higher altitudes.




Figure 6: Overview of fitted functions, soil moisture memory, its uncertainty (refer to text for details), $\tilde{\kappa}_n = \frac{\sigma_{w_n, y}}{\sigma_{P_n, y t_{lag}}}$ and $\rho(w_n, P_n)$ (as described in Section 4.4.1) for all catchments going from high soil moisture memory (top row) to low soil moisture memory (bottom row). In the left hand side column the red curves correspond to the fitted runoff ratio functions, the black lines show the fitted ET ratio functions and the vertical blue lines denote the 5% and 95% quantile of all soil moisture values in the time frame between April and October. Soil moisture memory, its uncertainty, $\tilde{\kappa}_n$ and $\rho(w_n, P_n)$ are computed for all months between April and October and for lag times between 5 and 40 days.




Figure 7: Continuation of Figure 6.




Figure 8: Correlations of fitted model parameters (listed in Table 1) at all catchments with respective soil moisture memory at a lag of 30 days. Dark gray corresponds to negative correlations, light gray indicates positive correlations. Hatching indicates correlations that are not significant on the 5% level (two-sided t-test).




Figure 9: The top row displays the soil moisture memories of lag 30 days of all 13 catchments plotted against altitude, CTI and dryness index including a least-squares fit and explained fraction of variance. The same is shown in the lower row for the ratio $\tilde{\kappa}_n = \frac{\sigma_{w_{n,y}}}{\sigma_{P_{n,y}t_{lag}}}$, also for a lag time of 30 days.